

Drug Interaction Report

5 potential interactions found for the following 4 drugs:

- **Benadryl** (diphenhydramine)
- **dexamethasone**
- **Haldol** (haloperidol)
- **lorazepam**

Add another drug

Interactions between your drugs

Moderate **haloperidol** ⇌ **LORazepam**
Applies to: Haldol (haloperidol), lorazepam

Using haloperidol together with LORazepam may increase side effects such as dizziness, drowsiness, confusion, and difficulty concentrating. Some people, especially the elderly, may also experience impairment in thinking, judgment, and motor coordination. You should avoid or limit the use of alcohol while being treated with these medications. Also avoid activities requiring mental alertness such as driving or operating hazardous machinery until you know how the medications affect you. Talk to your doctor if you have any questions or concerns. It is important to tell your doctor about all other medications you use, including vitamins and herbs. Do not stop using any medications without first talking to your doctor.

Switch to professional interaction data

Moderate **haloperidol** ⇌ **diphenhydrAMINE**
Applies to: Haldol (haloperidol), Benadryl (diphenhydramine)

Before using haloperidol, tell your doctor if you are also on diphenhydrAMINE. You may need dose adjustments or special tests in order to safely use both medications together. You should notify your doctor if you have signs of bladder problems, dry mouth, stomach pain, fever, blurred vision, confusion, dizziness, or reduced heart rate. You should avoid driving until you know how these medications will affect you. It is important to tell your doctor about all other medications you use, including vitamins and herbs. Do not stop using your medications without first talking to your doctor.

Switch to professional interaction data

Moderate

LORazepam ⇌ diphenhydrAMINE

Applies to: lorazepam, Benadryl (diphenhydramine)

Using LORazepam together with diphenhydrAMINE may increase side effects such as dizziness, drowsiness, confusion, and difficulty concentrating. Some people, especially the elderly, may also experience impairment in thinking, judgment, and motor coordination. You should avoid or limit the use of alcohol while being treated with these medications. Also avoid activities requiring mental alertness such as driving or operating hazardous machinery until you know how the medications affect you. Talk to your doctor if you have any questions or concerns. It is important to tell your doctor about all other medications you use, including vitamins and herbs. Do not stop using any medications without first talking to your doctor.

Switch to professional interaction data

No other interactions were found between your selected drugs. However, this does not necessarily mean no other interactions exist. Always consult your healthcare provider.

Drug and food interactions

Moderate

haloperidol ⇌ food

Applies to: Haldol (haloperidol)

Alcohol can increase the nervous system side effects of haloperidol such as dizziness, drowsiness, and difficulty concentrating. Some people may also experience impairment in thinking and judgment. You should avoid or limit the use of alcohol while being treated with haloperidol. Do not use more than the recommended dose of haloperidol, and avoid activities requiring mental alertness such as driving or operating hazardous machinery until you know how the medication affects you. Talk to your doctor or pharmacist if you have any questions or concerns.

Switch to professional interaction data

Moderate

LORazepam ⇌ food

Applies to: lorazepam

Alcohol can increase the nervous system side effects of LORazepam such as dizziness, drowsiness, and difficulty concentrating. Some people may also experience impairment in thinking and judgment. You should avoid or limit the use of alcohol while being treated with LORazepam. Do not use more than the recommended dose of LORazepam, and avoid activities requiring mental alertness such as driving

or operating hazardous machinery until you know how the medication affects you. Talk to your doctor or pharmacist if you have any questions or concerns.

Switch to professional interaction data

Therapeutic duplication warnings

No warnings were found for your selected drugs.

Therapeutic duplication warnings are only returned when drugs within the same group exceed the recommended therapeutic duplication maximum.

Report options



Share by QR Code

Drug Interaction Classification

These classifications are only a guideline. The relevance of a particular drug interaction to a specific individual is difficult to determine. Always consult your healthcare provider before starting or stopping any medication.

Major Highly clinically significant. Avoid combinations; the risk of the interaction outweighs the benefit.

Moderate Moderately clinically significant. Usually avoid combinations; use it only under special circumstances.

Minor Minimally clinically significant. Minimize risk; assess risk and consider an alternative drug, take steps to circumvent the interaction risk and/or institute a monitoring plan.

Unknown No interaction information available.

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.